




talking sense about sex

Last updated April 2016
Copyright FPA
www.fpa.org.uk/contraception

Methods with no user failure – they do not depend on you remembering to take or use them




Contraceptive injection

What is it? • An injection of progestogen.

Effectiveness • Perfect use: over 99%.
• Typical use: around 94%.

Advantage • Lasts for eight or 13 weeks – you don't have to think about contraception during this time.

Disadvantage • Can't be removed from the body so side effects may continue while it works and for some time afterwards.




Implant

• A small, flexible rod put under the skin of the upper arm releases progestogen.

• Perfect use: over 99%.
• Typical use: over 99%.

• Works for three years but can be taken out sooner.

• It requires a small procedure to fit and remove it.




Intrauterine system (IUS)

• A small, T-shaped, progestogen-releasing, plastic device is put into the uterus.

• Perfect use: over 99%.
• Typical use: over 99%.

• Works for 3–5 years but can be taken out sooner. Periods often become lighter, shorter and less painful.

• Irregular bleeding or spotting is common in the first six months.



Intrauterine device (IUD)

• A small plastic and copper device is put into the uterus.

• Perfect use: over 99%.
• Typical use: over 99%.

• Can stay in 5–10 years depending on type but can be taken out sooner.

• Periods may be heavier, longer or more painful.

Female and male sterilisation

• The fallopian tubes in women or the tubes carrying sperm in men (vas deferens) are cut, sealed or blocked.


• Failure rate is about 1 in 200 or 1 in 500 for females (depending on method), and 1 in 2,000 for males.

• Sterilisation is permanent with no long or short-term serious side effects.

• Should not be chosen if in any doubt about having children in the future.




Methods with user failure – you have to use and think about them regularly or each time you have sex




Contraceptive patch

What is it? • A small patch stuck to the skin releases estrogen and progestogen.




Contraceptive vaginal ring

• A small, flexible, plastic ring put into the vagina releases estrogen and progestogen.




Combined pill (COC)

• A pill containing estrogen and progestogen, taken orally.




Progestogen-only pill (POP)

• A pill containing progestogen, taken orally.




Male condom

• A very thin latex (rubber) polyurethane (plastic) or synthetic rubber sheath, put over the erect penis.




Female condom

• Soft, thin polyurethane sheath that loosely lines the vagina and covers the area just outside.



Diaphragm/cap with spermicide

• A flexible latex (rubber) or silicone device, used with spermicide, is put into the vagina to cover the cervix.



Natural family planning

• Fertile and infertile times of the menstrual cycle are identified by noting different fertility indicators.

PERFECT USE MEANS USING THE METHOD CORRECTLY EVERY TIME. TYPICAL USE IS WHEN YOU DON'T ALWAYS USE THE METHOD CORRECTLY.

Effectiveness	Contraceptive patch	Contraceptive vaginal ring	Combined pill (COC)	Progestogen-only pill (POP)	Male condom	Female condom	Diaphragm/cap with spermicide	Natural family planning
Perfect use: over 99%	Perfect use: over 99%	Perfect use: over 99%	Perfect use: over 99%	Perfect use: over 99%	Perfect use: 98%	Perfect use: 95%	Perfect use: 92–96%	Perfect use: up to 99%
Typical use: around 91%	Typical use: around 91%	Typical use: around 91%	Typical use: around 91%	Typical use: around 91%	Typical use: around 82%	Typical use: around 79%	Typical use: 71–88%	Typical use: around 76%
Advantage	• Can make bleeds regular, lighter and less painful.	• One ring stays in for three weeks – you don't have to think about contraception every day.	• Often reduces bleeding and period pain, and may help with premenstrual symptoms.	• Can be used by women who smoke and are over 35, or those who are breastfeeding.	Condoms are the best way to help protect yourself against sexually transmitted infections.		• Can be put in any time before sex.	• No physical side-effects, and can be used to plan as well as prevent pregnancy.
Disadvantage	• May be seen and can cause skin irritation.	• You must be comfortable with inserting and removing it.	• Missing pills, vomiting or severe diarrhoea can make it less effective.	• Late pills, vomiting or severe diarrhoea can make it less effective.	• May slip off or split if not used correctly or if wrong size or shape.	• Not as widely available as male condoms.	• You need to use the right size. If you have sex again extra spermicide is needed.	• Need to avoid sex or use a condom at fertile times of the cycle.